



UNITED WATER CONSERVATION DISTRICT

"Conserving Water Since 1927"

**Fish Passage Monitoring and Studies
Freeman Diversion Facility
Santa Clara River, Ventura County, California**



**Annual Report
2014 Monitoring Season**

This page left blank intentionally



UNITED WATER CONSERVATION DISTRICT

"Conserving Water Since 1927"

Fish Passage Monitoring and Studies
Freeman Diversion Facility
Santa Clara River, Ventura County, California

Annual Report
2014 Monitoring Season

Prepared by:

Steve Howard
and
Mike Booth

UNITED WATER CONSERVATION DISTRICT

106 North Eighth Street

Santa Paula, California 93060

2014

TABLE OF CONTENTS

	Page
List of Tables	iv
List of Figures	v
List of Photographs	vi
Executive Summary	vii
1.0 Introduction	1-1
1.1 Freeman Diversion Facility	1-1
1.2 Fish Species Composition	1-1
1.3 Environmental Setting	1-3
1.3.1 Rainfall.....	1-3
1.4 Regulatory Status (Steelhead)	1-4
2.0 Downstream Migrant Fish Passage	2-1
2.1 Introduction.....	2-1
2.2 Methods.....	2-1
2.2.1 Downstream Migrant Trap Checks	2-1
2.2.2 Smolt Length Measurements	2-5
2.3 Results	2-5
2.3.1 Downstream Migrant Trap Checks	2-5
2.3.2 Smolt Length Measurements	2-6
3.0 Upstream Migrant Fish Passage	3-1
3.1 Introduction.....	3-1
3.2 Methods.....	3-1

3.2.1	Bypass Flow Monitoring and Ladder Operation	3-2
3.2.2	Video Migration Monitoring	3-4
3.2.3	Ladder Shutdowns	3-5
3.2.4	DIDSON Migration Monitoring	3-6
3.3	Results	3-6
3.3.1	Bypass Flow Monitoring and Ladder Operation	3-6
3.3.2	Video Migration Observation.....	3-6
3.3.3	Ladder Shutdowns	3-6
3.3.4	DIDSON Migration Monitoring	3-6
4.0	Fish Stranding Surveys	4-1
4.1	Introduction.....	4-1
4.2	Methods.....	4-2
4.2.1	Fish Screen Bay Stranding Surveys	4-2
4.2.2	Lower Santa Clara River Stranding Surveys	4-2
4.2.3	Fish Ladder Stranding Surveys.....	4-3
4.3	Results	4-3
4.3.1	Fish Screen Bay Stranding Surveys	4-3
4.3.2	Lower Santa Clara River Stranding Surveys	4-3
4.3.3	Fish Ladder Stranding Surveys.....	4-4
5.0	Water Quality	5-1
5.1	Introduction.....	5-1
5.2	Methods.....	5-1
5.3	Results	5-2
5.1.1	Freeman Fish Trap Bay	5-2
5.1.2	Freeman Fish Screen Bay	5-2

5.1.3	Santa Clara River at VCWPD Project.....	5-3
5.1.4	Santa Clara River at VCWPD Project side channel	5-3
5.1.5	Santa Paula Creek upstream of Harvey Diversion.....	5-4
5.1.6	Santa Paula Creek at Steckel Park.....	5-4
5.1.7	Santa Paula Creek Directly Downstream of Sisar Creek Confluence	5-4
5.1.8	Sespe Creek at Grand Avenue	5-4
5.1.9	Piru Creek at Santa Felicia Dam.....	5-5
5.1.10	Piru Creek Upstream of Santa Clara River Confluence	5-5
5.4	Discussion	5-6
6.0	References.....	6-1
Appendix A. Tables and Figures		
Appendix B. Photos		

LIST OF TABLES

Page

Table 1. Freeman Diversion operations and monitoring results.....	Appendix A
Table 2. Santa Clara River fish species	Appendix A
Table 3. Fish acclimation schedule.....	2-4
Table 4. Freeman Diversion fish monitoring species results.....	Appendix A
Table 5. Santa Clara River reptile and amphibian species	Appendix A
Table 6. Temperature monitoring sites	Appendix A
Table 7. Water temperature statistics by monitoring site	Appendix A

LIST OF FIGURES

	Page
Figure 1. Schematic of Freeman Diversion Facility	Appendix A
Figure 2. Santa Paula rainfall	Appendix A
Figure 3. Steelhead monitoring results	Appendix A
Figure 4. Freeman bypass flow monitoring sites	Appendix A
Figure 5. Water temperature monitoring sites map.....	Appendix A
Figure 6. Freeman fish trap bay water temperature	Appendix A
Figure 7. Freeman fish screen bay water temperature	Appendix A
Figure 8. Santa Clara River at Ventura county bank stabilization project water temperature	Appendix A
Figure 9. Santa Clara River at Ventura county bank stabilization project side channel water temperature.....	Appendix A
Figure 10. Santa Paula Creek upstream of Harvey Diversion water temperature ..	Appendix A
Figure 11. Santa Paula Creek at Steckel Park water temperature	Appendix A
Figure 12. Sespe Creek at Grand Avenue water temperature	Appendix A
Figure 13. Piru Creek at Santa Felicia Dam level logger water temperature	Appendix A
Figure 14. Piru Creek at Santa Felicia Dam water temperature.....	Appendix A
Figure 15. Piru Creek at Santa Clara River confluence water temperature	Appendix A

LIST OF PHOTOGRAPHS

	Page
Photo 1. Santa Clara River estuary release site open	Appendix B
Photo 2. Santa Clara River estuary release site closed	Appendix B
Photo 3. Santa Clara River steelhead smolt	Appendix B
Photo 4. Santa Clara River resident rainbow trout	Appendix B
Photo 5. Upstream migration monitoring infrared scanners	Appendix B
Photo 6. Upstream migration monitoring surveillance system.	Appendix B
Photo 7. Fish screen bay stranding survey	Appendix B
Photo 8. Fish trap bay water temperature monitoring site	Appendix B
Photo 9. Fish screen bay water temperature monitoring site	Appendix B
Photo 10. VCWPD project water temperature monitoring site	Appendix B
Photo 11. VCWPD project side channel water temperature monitoring site	Appendix B
Photo 12. Santa Paula Creek upstream of Harvey Diversion water temperature monitoring site	Appendix B
Photo 13. Santa Paula Creek at Steckel Park water temperature monitoring site 2010	Appendix B
Photo 14. Santa Paula Creek at Steckel Park water temperature monitoring site 2014	Appendix B
Photo 15. Sespe Creek at Grand Avenue water temperature monitoring site	Appendix B
Photo 16. Piru Creek downstream of Santa Felicia Dam water temperature monitoring site	Appendix B
Photo 17. Piru Creek at Santa Clara River confluence water temperature monitoring site	Appendix B
Freeman Diversion Panoramic Photos	Appendix B

The water year of 2014 was considered a dry year. Rainfall in wet years can exceed 40 inches in the lower coastal plain and even more in the mountain areas. Rainfall in dry years can range from close to zero to a few inches. A total of 6.15 inches of rain was measured at the Santa Paula Wilson Ranch gauge (Station 245B) located in Santa Paula during the 2014 water year. A water year spans from October 1 of the previous year to September 30 of the current year. The highest monthly rain total in 2014 was 3.01 inches in February.

The sandbar at the Santa Clara River Estuary (SCRE) was opened for a short duration in the middle and end of February (February 11-14 and 28) and in March (March 1-19) during the only significant, although mild, storm of the year. Smolt trapping at the Freeman Diversion started on March 5, 2014 and ended on April 30, 2014 due to extremely low flows. A total of 11 steelhead smolts, 4 young-of-the-year, and 4 resident rainbow trout, were captured in the fish trap in 2014. No adult steelhead passed through the fish ladder during the migration season. No Pacific lamprey were observed during the 2014 migration season. Due to low flows and minimal rain, the fish ladder was only operated from March 2 to March 7, 2014.

Table 1, Appendix A summarizes the daily fish ladder operations as well as the *O. mykiss* and physical data collected during monitoring activities in 2014.

Water temperature monitoring in 2014 occurred in the mainstem Santa Clara River, Santa Paula Creek, Sespe Creek, and lower Piru Creek. Unfortunately, data uploading did not occur in the estuary but we will attempt to locate and upload the estuary temperature loggers if the estuary breaches in the 2015 water year.

1.1 FREEMAN DIVERSION FACILITY

The Freeman Diversion Facility was constructed in 1991 and is located approximately 10.7 miles upstream from the Pacific Ocean. The main purpose of the facility is to divert surface water from the Santa Clara River to conserve groundwater resources in the Oxnard plain through percolation to the groundwater aquifer. The facility is comprised of a concrete structure, a Denil fishway (fish ladder), a fish screen bay, a downstream migrant trap, fish trap bypass pipe, and various canals and spreading grounds (Figure 1, Appendix A). The concrete structure is a complete barrier to steelhead and potential barrier to Pacific lamprey upstream migration. The structure is only a potential barrier to lamprey because this species can traverse vertical obstructions if water is flowing over the surface and there are no sharp angles. The fish ladder was constructed to facilitate anadromous migration over the Freeman Diversion Structure. The fish screen bay is located directly downstream of where flow enters the facility and its function is to keep fish out of the canals and spreading grounds and to direct fish to the downstream migrant trap or to a fish trap bypass pipe. The fish trap bypass pipe is located at the end of the fish screen bay and can be used to direct fish back to the river when there is sufficient flow to allow for volitional migration to the estuary.

1.2 FISH SPECIES COMPOSITION

The Santa Clara River is home to two native, anadromous fish species, the southern California steelhead trout (*Oncorhynchus mykiss irideus*) and the Pacific lamprey (*Entosphenus tridentatus*). Steelhead and resident rainbow trout are known collectively as coastal rainbow trout. Steelhead is the common name for anadromous coastal rainbow trout. The term “anadromy” describes a life cycle or life history trait where species live as adults in the ocean and return to freshwater to spawn. Resident coastal rainbow trout live their entire lives in freshwater. Both resident as well as anadromous coastal rainbow trout exist in the Santa Clara River. Evidence from data collected at the Freeman Diversion

suggests that resident rainbow trout can produce progeny that will migrate to the ocean. A steelhead that migrates from freshwater where it was born to the ocean between the ages of 1 to 3 years is referred to as a smolt. The term smolt reflects the physical and physiological changes coastal rainbow trout experience when preparing for life in saltwater. A kelt is an adult steelhead that enters freshwater to spawn and later the same year (or the following year) returns to the ocean. Unlike Pacific salmon, steelhead do not necessarily die after spawning and can repeat several migration cycles to the ocean and again to freshwater to spawn. Southern steelhead are federally listed as endangered.

Pacific lamprey are strictly anadromous and cannot persist in freshwater alone. Juvenile lamprey, which live in freshwater for up to seven years before migrating to the ocean, are referred to as ammocoetes. Ammocoetes have no eyes and burrow in sandy sediments or live under larger substrates and debris. A downstream migrant Pacific lamprey is referred to as a macrophthalmia. The term macrophthalmia, similar to smolt, reflects the physical and physiological changes juvenile Pacific lamprey experience when preparing for life in saltwater, including the development of eyes. Pacific lamprey currently have no federal protection, but lamprey numbers have dropped precipitously since the early 2000's in the Santa Clara River and in many drainages on the west coast of the United States (Swift and Howard 2009).

There are two additional native fish species in the Santa Clara River, the federally-endangered tidewater goby (*Eucyclogobius newberryi*), which lives in the estuary and the threespine stickleback (*Gasterosteus aculeatus*). The threespine stickleback is comprised of two sub-species in the Santa Clara River, the partially-armored (*Gasterosteus aculeatus microcephalus*) and unarmored threespine stickleback (*Gasterosteus aculeatus williamsoni*). The partially-armored stickleback, which is plentiful and has no federal protection, exists in the Ventura county reach and the unarmored threespine stickleback exists in the Los Angeles county reach of the Santa Clara River. The unarmored threespine stickleback sub-species is a federal and California endangered species and a California fully-protected species. Additional fish species known to occur in the Santa Clara River include: Arroyo chub (*Gila orcutti*), Santa Ana sucker (*Catostomus santaanae*), Owens sucker (*Catostomus fumeiventris*), Santa Ana-Owens sucker hybrids (*C. santaanae* x *fumeiventris*), largemouth bass

(*Micropterus salmoides*), green sunfish (*Lepomis cyanellus*), bluegill (*Lepomis macrochirus*), brown bullhead (*Ameiurus nebulosus*), black bullhead (*Ameiurus melas*), prickly sculpin (*Cottus asper*), fathead minnow (*Pimephales promelas*), Mississippi (inland) silverside (*Menidia audens*), threadfin shad (*Dorosoma petenense*), common carp (*Cyprinus carpio*), goldfish (*Carassius auratus*), crappie (*Pomoxis spp.*), mosquitofish (*Gambusia affinis*), and shimofuri goby (*Tridentiger bifasciatus*). This is the first time shimofuri gobies have been documented in the mainstem Santa Clara River outside of the Piru Creek watershed. This species was first documented in the Piru Creek watershed within Pyramid Lake in 1990 (Swift et al. 1993). Swift et al. described the occurrence in Pyramid Lake as chameleon goby (*Tridentiger trigonocephalus*) but later recognized it as shimofuri goby (C.C. Swift pers comm). This species was observed by Steve Howard and Mike Booth in Piru Creek directly below Santa Felicia Dam in December 2013 and then in the Freeman Diversion fish trap in March 2014. See Table 2, Appendix A.

1.3 ENVIRONMENTAL SETTING

The Santa Clara River is the second largest watershed in southern California south of Point Conception, next to the Santa Ana River, and drains an area of approximately 1600 square miles. Its headwaters originate on the north slope of the San Gabriel Mountains near Acton, California in Los Angeles County and the river flows approximately 116 miles from east to west to its estuary in the City of Ventura. The Santa Clara River and its tributaries have high annual flow variability, from extreme flood events to multi-year droughts.

1.3.1 RAINFALL

During the 2014 rain season (10/1/2013-9/30/2014), Santa Paula had 35.4% of normal rainfall totaling 6.15 inches (Ventura Watershed Protection District website, Santa Paula-Wilson Ranch 245B). The most rainfall in a 24 hour period was 1.62 inches on February 28, 2014 and the second highest was 1.13 inches on both February 27 and March 1, 2014. The highest monthly rainfall was 3.01 inches in February. (Figure 2, Appendix A). Southern California can have cold or warm winters depending on ocean currents and other factors. Cold winters, tend to produce moderate storms and snow on local mountains. Warm winters, such as 2005, have storms that produce large amounts of rain over a short period of

time, typically resulting in flood conditions. Cold winters can be beneficial to *O. mykiss* because of sustained cold water flows from snow melt sometimes occurring well into the summer months. Warm winters can have negative effects on *O. mykiss* due to redd scouring but can also be beneficial when high flows remove excess vegetation (both exotic and native), scour and transport fine sediments, deposit clean gravels, and provide extended migration opportunities.

Tributaries

The major tributaries of the Santa Clara River include Santa Paula Creek, Sespe Creek, and Piru Creek. Santa Paula Creek has two fish passage facilities located within the first four miles of the lower creek (USACE and Canyon Irrigation fish ladders). These facilities have numerous physical and operational problems that result in restricted upstream passage either intermittently or completely depending on maintenance or damage in any given year. Sespe Creek is free-flowing, but illegal road crossings within the lower river have had the potential to block both upstream and downstream passage at certain flows in the past. Piru Creek has two major dams (Santa Felicia Dam and Pyramid Dam) that do not include fish passage facilities and are complete barriers to upstream fish passage and intermittent barriers to downstream passage (fish can move downstream during spills over dams).

Santa Paula and Sespe Creeks are considered to be the primary steelhead-bearing drainages in the Santa Clara River watershed and are the closest major tributaries to the ocean. These tributaries did not have surface flows connecting to the Santa Clara River during much of the 2014 migration season. It is unknown when steelhead smolts emigrate from the tributaries to the mainstem Santa Clara River. Currently, it is assumed that smolts emigrate from the tributaries following storm pulses based on data collected in the Santa Ynez River (Tim Robinson, Cachuma Operations and Maintenance Board, pers. comm.). Steelhead smolts are typically observed at the Freeman Diversion from January to June but the majority of these fish are observed in March, April, and May.

1.4 REGULATORY STATUS (STEELHEAD)

NOAA Fisheries, otherwise known as the National Marine Fisheries Service

(NMFS), listed the southern California steelhead, *Oncorhynchus mykiss irideus*, as endangered in 1997 (NMFS 1997) under the Endangered Species Act (ESA) of 1973. Steelhead were organized into stocks (i.e., groups) of evolutionary significant units (ESU) that represented groupings considered to be substantially reproductively isolated from other steelhead stocks and an important part of the evolutionary legacy of the species. Currently, the southern California steelhead ESU includes populations from the Santa Maria River in San Luis Obispo County south to the US/Mexican border in San Diego County (NMFS 2003). NOAA Fisheries later recognized the anadromous life history form of *O. mykiss* as a distinct population segment (DPS) under the ESA (NMFS 2006). The DPS policy differs from the ESU by delineating a group of organisms by “marked separation” rather than “substantial reproductive isolation”. In the case of *O. mykiss* of the southern California steelhead ESU, this marked separation between the two life history forms was considered valid because of physical, physiological, ecological, and behavioral factors related to its anadromous life history characteristics. Both resident and anadromous *O. mykiss* (where the two forms co-occur, below complete barriers, and are not reproductively isolated) are still part of the ESU; however, the anadromous *O. mykiss* (steelhead) are now part of a smaller subset identified as the southern California steelhead DPS (CMWD 2008).

The remainder of this report summarizes the monitoring methods and results at the United Water Conservation District (hereafter, District) Freeman Diversion Fish Passage Facility on the Santa Clara River in 2014.

2.1 INTRODUCTION

In southern California, steelhead and Pacific lamprey migrate downstream from their natal streams to the Pacific Ocean in the spring. Data collected from 1991 to 2014 at the Freeman Diversion indicate that the majority of downstream migration occurs in March, April, and May, although migration can occur from January through June and during 2009 migration was observed into July. Migration typically occurs when flows in the Santa Clara River are rapidly receding. When this occurs, it is necessary to trap all downstream migrant steelhead and Pacific lamprey and relocate them to the estuary or other appropriate habitats (based on individual life stages) to prevent fish stranding in dewatered sections of the river below the Freeman Diversion.

The primary objective for trapping and transporting downstream migrants is to avoid impacts to steelhead smolts, kelts, and lamprey macrophthalmia during their downstream migration to the Pacific Ocean when there is not sufficient flow in the lower river. This reduction in flow could be natural and/or the result of diversions at the Freeman Diversion and by other water users in the watershed. Another important objective is to gather data regarding anadromous downstream migration in the Santa Clara River. Since there is a lack of knowledge regarding steelhead in southern California, these data can be useful during the steelhead recovery planning process and for managers of anadromous fishes in the Santa Clara River as well as the region as a whole. Additionally, trapping activities aid in monitoring fish movement and assemblages within the Santa Clara River and can potentially prevent stranding and predation when conditions are not favorable in the lower river due to natural conditions as well as diversions occurring at the Freeman Diversion and elsewhere in the watershed.

2.2 METHODS

2.2.1 DOWNSTREAM MIGRANT TRAP CHECKS

Trapping occurred from March 5 to April 30, 2014. Annual trapping typically starts

in January, but extremely low flows precluded trapping until the first storm occurred at the end of February. Fish relocation is triggered when there is insufficient flow in the lower river based on depth criteria at critical riffles. The current smolt target flow is 120 cfs measured approximately 5.5 mi downstream of the Freeman Diversion within a sandy glide habitat unit in the lower Santa Clara River. This target flow threshold was developed from an analysis of depth, velocity and wetted width taken at a range of flows within long, sandy glides. These results were used to develop a smolt target flow (120 cfs) where more than 50% of the wetted stream channel was greater than 0.5 feet deep. Trapping and transporting commenced when flow receded to 80 cfs at the sandy glide flow measurement site. A trapping trigger or critical passage flow of 80 cfs was developed because only a narrow width (10 % or less of the wetted stream channel) was equal to or greater than 0.5 feet deep during the depth, velocity and wetted width study. A smolt radio telemetry study is in development to address uncertainties regarding smolt migration behavior and rate of migration in the lower Santa Clara River. This study is planned to be implemented in United's Multi-Species Habitat Conservation Plan (MSHCP). The primary objective of this study will be to evaluate smolt migration rates and success at various flows to inform and potentially modify the smolt target flows and trapping operations.

Downstream migrant steelhead smolts, kelts, Pacific lamprey macrophthalmia, and other fish entering the fish screen bay within the diversion facility were prevented from entering the diversion canal by a self-cleaning, 1.75 mm (0.07 inch) mesh wedge wire screen which directs fish to a downstream migrant fish trap. Steelhead smolts trapped at the facility were transported in aerated coolers by truck to the Santa Clara River Estuary when flows were less than 80 cfs at the critical riffle, or released in the Santa Clara River below the Freeman Diversion when migration flows were being released and greater than 80 cfs. No steelhead adults, kelts or Pacific lamprey adults were observed or captured in 2014. No lamprey macrophthalmia or ammocoetes were observed or trapped. Resident coastal rainbow trout not exhibiting smolting characteristics were transported, depending on flow conditions, to the Santa Clara River, Santa Paula Creek, or Sespe Creek. Currently, the Santa Clara River above Santa Paula is the preferred relocation site for resident coastal rainbow trout and lamprey ammocoetes. All other native aquatic species were returned to the river upstream of the diversion. Non-native and exotic aquatic species were removed from the

river unless they were considered special status species in neighboring watersheds (e.g., Santa Ana sucker). Genetic research is currently in progress by researchers at the United States Geological Survey to evaluate whether Santa Ana sucker are native to the Santa Clara River. It has been speculated that this species was introduced based on a lack of observations or records in the early 1900's.

The downstream migrant trap consists of 3/16-inch mesh metal screens. Flow enters through a weir gate with an opening that directs fish and other aquatic species into the trap from the fish screen bay. The trap is situated to keep all intercepted fish immersed in at least two feet of flowing water.

The trap was checked daily in the morning and all fish were removed from the trap with a dip net, counted and measured. Other trapped aquatic species were counted and documented during each trap check. All fisheries personnel were trained in species identification and handling. A species identification handbook drafted by Steve Howard (senior ecologist) and Mike Booth (associate ecologist) was also available.

Data Collection – Steelhead smolts were measured (fork length) to the nearest millimeter on a wet fish measuring board. Measurements typically took no more than 10 seconds per individual. Lamprey ammocoetes and macrophthmia, when observed, are measured to total length from head to tail. Water temperature was measured in the fish trap prior to handling fish and monitored in the transport cooler during transport. Fish were not transported at temperatures above 20°C and transport activities were performed early in the morning to minimize stress. General fish condition was assessed as well as the general degree of smoltification or smolt condition (presmolt or full smolt). Water temperature, dissolved oxygen, pH, conductivity, salinity and turbidity were measured at the relocation sites with a multi-parameter water quality meter. When possible, photos of collected and transported steelhead and lamprey were taken with a digital camera. All data were documented on standardized datasheets and transferred daily to an electronic database.

Fish Transportation – Fish were collected from the fish trap with 1/8th inch or smaller mesh dip nets, data were collected and compiled on datasheets and the fish were typically placed in 100 quart aerated coolers. No more than 25 smolts,

ammocoetes or macrophthalmia were placed in an individual cooler. No more than one adult steelhead or kelt was placed in individual coolers. If a low number of smolts or macrophthalmia were collected, these fish were placed in aerated buckets to minimize handling during release. Fish handling and transport time was generally no more than one hour.

Fish Transport Locations

Anadromous Fish

All anadromous downstream migrant fish (steelhead smolts and kelts, Pacific lamprey macrophthalmia) were transported from the Freeman Diversion to the Santa Clara River Estuary (Photo 1 Appendix B) when discharge in the Santa Clara was less than 80 cfs. When flows were greater than 80 cfs, fish were released in the Santa Clara River downstream of the Freeman Diversion. The specific estuary relocation site depended on the condition of the estuary. The estuary was monitored daily during the migration season to inform relocation activities. Generally fish were released to fresh water in the estuary that had at least 1 foot of depth and instream cover nearby. Areas of the estuary that were known to be low in oxygen were avoided.

Acclimation Schedule - Fish were acclimated to the receiving water using the schedule in Table 3.

Table 3 – Fish acclimation schedule.

Degree Differential (between cooler and receiving water)	Acclimation Minutes
0-2	10
3-5	20
6-7	30
8 and over*	40

* Fish transportation should not occur when the water temperature is too high ($>20^{\circ}\text{C}$). If this occurs the fish transport time schedule will be modified. If the estuary water temperature is over 23°C the lead fisheries biologist will be contacted before acclimating.

Non-Anadromous Fish

Non-anadromous fish (resident coastal rainbow trout and Pacific lamprey ammocoetes) were transported via aerated coolers to either the mainstem Santa Clara River, Sespe Creek, or Santa Paula Creek (Photos 10, 15, and 13 respectively, Appendix B). If conditions were not suitable for trout in the mainstem Santa Clara River, resident rainbow trout might be transported to Santa Paula Creek upstream of Steckel Park or Sespe Creek at Grand Avenue. All other native, non-anadromous fish and aquatic species were transported and released upstream of the Freeman Diversion Structure.

2.2.2 SMOLT LENGTH MEASUREMENTS

Trapped smolts were measured to fork length. Lengths were measured by placing the fish in a fish measurement cradle for no more than 10 seconds. The fish were not sedated. If an individual fish could not be measured in less than 10 seconds, the fish was placed back in an aerated cooler and not measured again.

2.3 RESULTS

2.3.1 DOWNSTREAM MIGRANT TRAP CHECKS

The downstream migrant trap was in operation from March 5 to April 30, 2014. Eleven smolts were captured during 2014 (Figure 3, Appendix A). A total of 1 resident coastal rainbow trout and 4 young-of-the-year (YOY) coastal rainbow trout were trapped and relocated during the 2014 migration season. Three additional resident coastal rainbow trout were collected in the fish screen bay (see section 4.3.1)

Additional fish collected in the Freeman downstream migrant trap included: partially armored threespine stickleback (N=608), Arroyo chub (N=20), Santa Ana sucker (N=6), Owens sucker (N=17), Santa Ana/Owens sucker hybrids (N=4), fathead minnow (N=57), common carp (N=10), green sunfish (N=6), brown bullhead (N=1), black bullhead (N=13), prickly sculpin (N=52), shimofuri

goby (N=9) and mosquitofish (N=4262) (Table 4, Appendix A). Sucker species were identified based on lip morphology to the extent possible, but the validity of this method is questionable. Red swamp crayfish were regularly captured in the trap (N=2813).

Amphibians and reptiles collected in the Freeman downstream migrant fish trap included: Western toad (N=6), bullfrog (N=242, mostly larvae), African clawed frog (N=39), tree or Pacific chorus frog (N=1), and Western pond turtle (N=1) (Table 4, Appendix A). The regional and regulatory status of amphibians and reptiles collected at the Freeman Diversion can be found in Table 5 (Appendix A). A healthy pond turtle population exists directly above and below the Freeman Diversion Structure.

2.3.2 SMOLT LENGTH MEASUREMENTS

A total of 11 smolts were measured. The average length of the 2014 smolts was 145.3 mm (standard deviation 35.9 mm).

3.1 INTRODUCTION

The Freeman Diversion Facility is equipped with a Denil fish ladder (fishway) that was constructed to facilitate steelhead and Pacific lamprey upstream migration over the concrete diversion structure. Although Denil fishways are not ideal for lamprey passage, the fish ladder did pass hundreds of lamprey in the 1990's (Chase 2001, Swift and Howard 2009). There is some uncertainty regarding how many steelhead have passed the diversion structure because of a previously inconsistent and inefficient monitoring program, as well as regulatory and extreme environmental constraints (flashy flows, high turbidity). Despite uncertainties in the data, a minimum of 11 wild adult steelhead and 2 hatchery steelhead have been observed at the diversion structure since it was constructed in 1991.

3.2 METHODS

In years with sufficient rainfall, upstream migrant monitoring is conducted to determine if adult steelhead use and effectively navigate the fish ladder. Currently, there is no active fish trap deployed within the fish ladder to monitor steelhead and lamprey upstream migration. In order to monitor steelhead upstream migration, the District installed a passive monitoring device that counts upstream migrants when they jump over a false weir and through an infrared (IR) scanning device while then being documented by an integrated video monitoring system. Lamprey likely migrate under a grated section of the passive monitoring device, where it is unlikely that they will be observed. The only other option for monitoring steelhead or lamprey upstream migration within the diversion facility is during fish ladder shutdowns or when flow is reduced within the fish ladder. Monitoring during hour-long shutdowns of the fish ladder only allows for intermittent observations during the migration season. It would benefit steelhead recovery efforts and increase knowledge regarding steelhead and Pacific lamprey migration if an active trap was installed within the fish ladder. A trap would temporarily capture all upstream migrants so that information could be

gathered regarding ladder efficiency, migration timing, fish condition, water quality at migration, as well as other parameters of interest.

3.2.1 BYPASS FLOW MONITORING AND LADDER OPERATION

The current low flow threshold for sufficient upstream migration is 160 cfs, measured at critical riffles in the lower Santa Clara River. This threshold was developed from the results of a steelhead migration instream flow study conducted by Thomas R. Payne and Associates (now Normandeau Associates) in the lower Santa Clara River (TRPA 2005). This study evaluated surface water depths at various flows to understand what minimum flows would be required for steelhead to successfully migrate from the ocean to the Freeman Diversion Fish Ladder.

During fish ladder operations, bypass flows are monitored at two sites between the Freeman Diversion and Highway 101 to ensure that a minimum of 160 cfs is maintained at the most downstream monitoring site located approximately 0.6 miles upstream of the Highway 101 Bridge. The two monitoring sites are located approximately 0.14 miles below the Freeman Diversion Structure and 5.5 miles downstream of the Freeman Diversion or 0.6 miles upstream of the 101 Bridge (Figure 4, Appendix A). The most downstream monitoring site is located near the end of the losing reach of the river where the highest percolation rates of surface water are believed to occur.

Flow is measured with a YSI/SonTek FlowTracker Acoustic Doppler Velocimeter. Flow measurements are conducted using USGS standards for measuring flow using acoustic Doppler technology in open channels (Rehmel 2007). The measurement site (critical riffle area) is located within a long and wide sandy glide. The numerous shallow glides found in the Santa Clara River potentially result in migration bottlenecks and delay for steelhead migration (S. Howard, personal observations of physical habitat). The actual critical riffles (typical area to measure shallow conditions) have narrower active channels (wetted width) than glide habitats, resulting in deeper, more suitable migration conditions compared to the wide, sandy glides.

The Freeman Diversion Fish Ladder operations are based on negotiations between NMFS biologists and the District hydrologist and biologists. These

operating criteria were revised in the United States Bureau of Reclamation's *Biological Assessment of the Operation of the Vern Freeman Diversion Dam and Fish Ladder, Santa Clara River, Ventura County, California* and later revised in the Proposed 2009 Interim Operations Plan and 2010 Smolt Bypass Flow Plan.

The current bypass flow operating criteria are as follows:

When United starts diverting water after a storm (turn-in), diversion rates are limited when total river flow is less than or equal to 750 cfs. When turning in at total river flow greater than 635 cfs and less than or equal to 750 cfs, only 30% of the remaining river can be diverted providing that the required bypass flows for steelhead are met. When turning in at total river flow less than or equal to 635 cfs, only 20% of the remaining river can be diverted providing that the required bypass flows for steelhead are met. These diversion restrictions apply only to turning-in procedures undertaken during the principal steelhead migration season (January through May) when total river discharge is less than or equal to 750 cfs.

January 1 – March 14 Bypass Flows

After a storm event triggers migration flows, the fish ladder and auxiliary flows will be operated on the following schedule: From January 1 to March 14 (adult bypass flows), bypass flows with a target of 160 cfs including a ramp-down schedule will be maintained for 18 days. If total river flow drops below 160 cfs at the critical riffle without diversions during the 18 day flow window, the total amount of flow released will be reduced by 1/3 each day for three days and reduced to 20 cfs on the fourth day. If an adult steelhead is detected in the fish ladder during ramp down, flows will be maintained at the current release discharge for 24 hours and then ramp down will proceed if no adult steelhead are detected. During the ramp down period, United may divert water not being released downstream.

March 15 – May 31 Bypass Flows

From March 15 to May 31, United may continuously maintain critical diversions of 40 cfs during bypass flow releases. From March 15 to May 31, following a storm, a bypass flow of 160 cfs will be maintained for 11 days then reduced to 120 cfs until this flow cannot be maintained. The smolt target flow of 120 cfs was developed from water depth surveys focused on providing downstream migrating

smolts with safe, volitional passage through the lower river. However, when flows at the critical riffle are predicted to reach a low critical flow of 80 cfs within 5 days, smolts will be trapped and transported. After flows reach 80 cfs, bypass flow will be reduced by 1/3 each day for three days and reduced to 20 cfs on the fourth day. For all ramp down operations, if natural flows decline faster than the specified ramping rates and it is impossible to maintain the prescribed 1/3 daily reduction in flow, then natural flows will be used for that 24 hour period.

Smolts may migrate downstream regardless of storm events, so United will provide bypass flows of 120 cfs for smolt migration from March 15 to May 31 while maintaining critical diversions of 40 cfs. If there is not sufficient water to provide smolt bypass flows of 120 cfs and maintain the critical diversion of 40 cfs, smolts will be trapped and transported when river flows are predicted to recede to 80 cfs with 40 cfs diverted within five days. When adult bypass flows are not in effect and smolts are captured in the fish trap outside of the smolt bypass flow period, they will be transported to the estuary.

3.2.2 VIDEO MIGRATION MONITORING

Upstream migration monitoring is conducted using a false weir, infrared scanners, and a computer-based surveillance system (Photo 5 and 6, Appendix B). The false weir creates a barrier within the ladder that forces upstream migrant steelhead to jump out of the water approximately six inches to traverse a small “fall or plunge” to continue upstream. Consequently, the migrating steelhead jump through infrared beams that detect the migration event. Four cameras capture video of each fish that negotiates the weir and the surveillance system records video of the events on a computer hard drive. The weir was designed with a bar screen at the bottom of the structure to allow Pacific lamprey to pass under the false weir. It is unclear if Pacific lamprey are able to pass through the bar screen due to a lack of data regarding approach water velocities at the screen. Approach velocities at the bar screen and potential monitoring alternatives will be evaluated during the next several years of operations.

Prior to the deployment of the current computer surveillance system in 2010, a DVR system with a single low resolution camera was used to monitor steelhead migration through the fish ladder. The new computer based system provides substantially greater resolution and multiple functions to detect movement.

During 2014, IR scanners were used to detect movement over the false weir. Although the surveillance software can detect movement, it is difficult to isolate detection of fish from the water moving in the background. When steelhead traverse the fish ladder and jump over the false weir, the computer based monitoring system documents the event with a flashing alarm on the computer screen. Alarms and associated video footage are reviewed daily. The computer system records continuously and there is sufficient hard disk storage to save video for up to 7 days, depending on the number of cameras in use. Once the hard drive is full, new data is saved over old data, starting at the beginning of the old file. Migration events are saved as short video files on the computer hard drive. The surveillance system receives digital video from two low-light, high resolution, above-water cameras placed upstream and downstream of the false weir, as well as an underwater camera placed downstream focused on the false weir. Three twenty-five watt fluorescent lights were used to illuminate the false weir at night.

The IR scanner device is equipped with sensors on both sides of the false weir with multiple closely-spaced infrared beams. When the beams are broken by a fish jumping over the weir, the IR scanner signals the surveillance system to record the event.

The video monitoring system is checked daily to ensure proper function and to evaluate the efficiency of the IR scanner device. Triggered alarms are reviewed and any necessary adjustments are made to the water levels, scanning devices, or recording equipment. Two motion detection cameras (2 of 4 cameras total) were placed at the weir after the 2012 monitoring season to provide views conducive to using motion detection in the surveillance software. Motion detection as well as infrared detection is now in place to increase the efficiency of the detection system.

3.2.3 LADDER SHUTDOWNS

A thorough examination of the entire fish ladder is conducted to check for the presence of any fish species during shutdown or dewatering of the ladder. The ladder is shut down for operational or maintenance reasons during high storm flows, channel flushes, and/or removal of debris from the weir.

3.2.4 DIDSON MIGRATION MONITORING

DIDSON monitoring did not occur in 2014 due to the short duration of bypass flows and the steep recession of river flow. Our efforts were focused on the challenges of providing the required bypass flows as the river rapidly receded.

3.3 RESULTS

3.3.1 BYPASS FLOW MONITORING AND LADDER OPERATION

Bypass flows occurred following the storm on February 28 and ended when the required bypass flows could not be maintained and following ramp-down procedures on March 8. The fish ladder was in operation from March 2 to March 7, 2014. No adult steelhead or Pacific lamprey were observed in 2014.

3.3.2 VIDEO MIGRATION OBSERVATION

The surveillance system was operated continually from March 2 to March 7, 2014 during the operation of the fish ladder. Throughout this period the alarm trigger system was monitored and checked daily.

3.3.3 LADDER SHUTDOWNS

The fish ladder was shut down only once in 2014 at the end of ladder operations on March 7.

3.3.4 DIDSON MIGRATION MONITORING

No DIDSON monitoring occurred in 2014.

4.1 INTRODUCTION

During high flow events when the river is highly turbid from elevated concentrations of total suspended solids (over 3000 ntu's), District operations staff "turn-out" all river flows from the facility resulting in the impound upstream of the diversion structure becoming dewatered or extremely shallow. When water is turned out of the facility, head gates are closed to retain sufficient depth in the fish screen bay to avoid potential predation of fish by birds until fisheries staff can survey for stranded fish. During all "turn-out" events when the fish ladder is in operation, the fish ladder must be shut down and inspected for potential stranded fish. When the fish ladder is shut down it slowly dewateres and this can result in fish becoming stranded in the fish ladder.

Additional operations and maintenance activities include "flushes" where District operations staff temporarily "turn-out" all river flows from the facility to maintain the active channel toward the facility headworks or to conduct maintenance of the canal gates and screens. During a flush, the river will pass through the flushing channel (also known as bypass channel), carrying away material deposited upstream of the intake. As the flush continues, the river continues to scour sediment or headcut for a distance back from the intake, up to several hundred feet. Flushing is necessary to keep both the diversion and the fish ladder in operation. When flushes occur, head gates are closed to retain sufficient depth in the fish screen bay to avoid potential predation of fish by birds until fisheries staff can survey for stranded fish.

The primary objective of fish stranding surveys is to rescue any fish that become stranded when diversion operations cause river flow to rapidly diminish downstream of the Freeman Diversion in the Santa Clara River and within the fish screen bay and fish ladder during dewatering operations.

4.2 METHODS

4.2.1 FISH SCREEN BAY STRANDING SURVEYS

During turn-out and flushing events, the fish screen bay was thoroughly examined for the presence of steelhead, Pacific lamprey, and other aquatic species. Two or more biologists surveyed the fish screen bay as it dewatered. These surveys were conducted by seining the entire area with either 1/8 or 1/4 inch mesh brailed seines, 4 feet deep and 10 to 20 ft long (Photo 7, Appendix B). The primary objective of these surveys was to capture and relocate steelhead and Pacific lamprey stranded during dewatering of the fish screen bay to appropriate habitats either in the estuary (smolts, kelts, macrophthalmia) or the Santa Clara River and associated tributaries (resident coastal rainbow trout and Pacific lamprey ammocoetes). Also, the fish screen bay was periodically dewatered opportunistically to look for steelhead and Pacific lamprey migrants that could potentially be delayed in the fish screen bay. All fish were transported via aerated coolers and non-native aquatic species were removed from the river. All data collected during stranding surveys were documented on standardized datasheets. Fish were transported utilizing methods described in Section 2.2 of this report.

4.2.2 LOWER SANTA CLARA RIVER STRANDING SURVEYS

In 2014, no surveys were required. In previous years, stranding surveys were conducted in the Santa Clara River below the Freeman Diversion when bypass flows were significantly reduced or when releases were stopped all together, due to turn-out or flushing events. These surveys were conducted by entering the non-wetted area of the floodplain via a Polaris Ranger all-terrain vehicle that was stocked with equipment necessary to collect, hold and transport stranded fish. Stranded fish were captured either with dip nets or 1/8 inch to 1/4 inch mesh brailed seines and placed in aerated coolers. In years when upstream migrant steelhead and Pacific lamprey are collected during these surveys they will be relocated upstream of the Freeman Diversion Structure. Downstream migrant steelhead (smolts) will be relocated to the Santa Clara River Estuary. In years when downstream migrant Pacific lamprey (macrophthalmia) are collected during stranding surveys they will be relocated to the estuary.

4.2.3 FISH LADDER STRANDING SURVEYS

Stranding surveys were conducted in the fish ladder during turn-out events. When the fish ladder is in operation and one of these events occurs, the fish ladder becomes dewatered and the potential exists for steelhead and Pacific lamprey to become stranded either between the Denil plates or in resting pools of the fish ladder. A biologist was present and conducted surveys when these operations occurred by walking the length of the ladder searching for any stranded fish as the ladder dewatered. Dip nets and buckets were used to rescue any stranded fish.

Fish ladder stranding surveys were also conducted during maintenance activities to visually survey for steelhead and Pacific lamprey that that might become stranded in the fish ladder during these activities. If a fish was observed in the Denil sections of the ladder it was captured with a dip net, placed in a bucket and relocated to an appropriate location based on its life stage.

4.3 RESULTS

4.3.1 FISH SCREEN BAY STRANDING SURVEYS

A total of one turn-out and zero flush events occurred during the 2014 water year. The turn-out events were due to high turbidity levels from the suspended sediment load, while the flush was performed for channel maintenance and personnel training. One fish screen bay stranding survey was conducted at the end of trapping operations due to receding river flows and diversions due to drought conditions. The one stranding survey yielded three resident trout (Table 4, Appendix A). The resident trout were released in the mainstem Santa Clara River at the Ventura County Watershed Protection District project near Santa Paula. These fish were included in the total trapping results section (see section 2.3.1).

4.3.2 LOWER SANTA CLARA RIVER STRANDING SURVEYS

No visual stranding surveys were required below the Freeman Diversion structure following turn-out events because bypass flows were released following the turn-out events which maintained surface flow in the river.

4.3.3 FISH LADDER STRANDING SURVEYS

The fish ladder was surveyed for stranded fish only once in 2014 when the ladder was shut down on March 7. A biologist walked the length of the fish ladder during this event as the ladder dewatered. No *O. mykiss* were observed stranded in the fish ladder during this survey.

5.1 INTRODUCTION

Water quality monitoring for 2014 was conducted to monitor water quality conditions that steelhead and Pacific lamprey are exposed to in various areas within the watershed. Water quality parameters such as dissolved oxygen, pH, conductivity, salinity, and turbidity were collected *in situ* at some sites. Water temperature monitoring was conducted in various locations in the watershed and water quality measurements were focused primarily in the Santa Clara River Estuary, Santa Paula Creek, Sespe Creek, and the main stem Santa Clara River where coastal rainbow trout were relocated during the 2014 monitoring season. Only water temperature measurement results are included in this report because other parameters were only measured during intermittent fish transport activities.

5.2 METHODS

Water quality monitoring in 2014 consisted of water temperature measurements utilizing Onset U22 Hobo® temperature loggers at various sites within the Santa Clara River watershed and in-situ water quality measurements taken during fish relocation activities using either a Hach Quanta or Horiba U-10 multi-parameter water quality meter. The majority of the temperature loggers were placed in pool habitats at mid-depth and programmed to take measurements every hour. The standardized data collection dates were October 1, 2013 through September 31, 2014.

Water temperature monitoring occurred at the following sites:

- Site 2: Freeman Fish Trap Bay
- Site 3: Freeman Fish Screen Bay
- Site 4: Santa Clara River at the Ventura County bank stabilization project
- Site 4A: Santa Clara River side channel at the Ventura County bank stabilization project
- Site 6B: Santa Paula Creek upstream of Harvey Diversion

- Site 7: Santa Paula Creek at Steckel Park
- Site 9B: Sespe Creek at Grand Avenue
- Site 17B: Piru Creek at Santa Felicia Dam
- Site 17C: Piru Creek at Santa Felicia Dam (Level Logger)
- Site P1: Piru Creek upstream of confluence with Santa Clara River

Some of the other site numbers are not presented here because the loggers were either lost due to high flows or conditions were such that access to the sites was not feasible. A map identifying each monitoring site can be found in Figure 5, Appendix A. Graphs depicting data at these eight sites are in Figures 6-15, Appendix A. Specific information regarding each site can be found in Table 6, Appendix A.

5.3 RESULTS

5.1.1 FREEMAN FISH TRAP BAY

The Santa Clara River Freeman Fish Trap Bay water temperature monitoring site was located within the fish trap bay at the Freeman Diversion in Saticoy. The logger was attached to a rung of a permanent ladder structure within the fish trap bay. The logger was attached with zip ties approximately 1.0 feet from the bottom of the bay (Photo 8, Appendix B). The objective at this location was to monitor water temperatures in the fish trap bay where the downstream migrant fish trap is located. Steelhead smolts and several other aquatic species were trapped at this location during the migration season (typically January-June). Temperature monitoring is critical at this location as this is where the fish remain until the daily trap check occurs. Occasionally resident and young-of-the-year coastal rainbow trout as well as steelhead kelts may also be found in the trap. The minimum water temperature collected at this site was 1.72 °C and the maximum temperature was 31.5 °C (Table 7, Appendix A). This site was dewatered during much of the season, which is reflected in the maximum and minimum values.

5.1.2 FREEMAN FISH SCREEN BAY

The Santa Clara River Freeman Fish Screen Bay water temperature monitoring site was located within the fish screen bay. The fish screen bay consists of a concrete channel where downstream migrants enter the diversion and migrate to the downstream migrant fish trap. The logger was located approximately 50 feet

downstream from the intake to the fish screen bay and attached to an eye hook 0.5 feet from the bottom along a concrete wall (Photo 9, Appendix B). The objective at this location was to monitor water temperatures in the main stem as river flow enters the Freeman Facility. Downstream migrant steelhead, Pacific lamprey and other aquatic species temporarily rear in the fish screen bay prior to entering the trap. The downstream migrant fish trap is typically taken out of operation between June and December and all fish that enter the facility rear in the fish screen bay until trapping operations commence in January. The minimum water temperature collected at this site was -0.17 °C and the maximum temperature was 47.1 °C (Table 7, Appendix A). This site was dewatered during much of the season, which is reflected in the maximum and minimum values.

5.1.3 SANTA CLARA RIVER AT VCWPD PROJECT

The Santa Clara River at the Ventura County Watershed Protection District (VCWPD) Project water temperature monitoring site was located approximately 100 feet downstream from the most downstream groin at the county bank stabilization project site off South Mountain Road in Santa Paula. The logger was attached to a fence post with zip ties 0.9 feet from the bottom (Photo 10, Appendix B) in a pool about 3.5 ft deep. The objective at this location was to monitor water temperatures in the main stem Santa Clara River between Sespe Creek and the Freeman Diversion. In the future we will be adding more temperature monitoring sites throughout the Santa Clara River to closely monitor temperatures during the steelhead and Pacific lamprey migration season. The minimum water temperature collected at this site was 10.4 °C and the maximum temperature was 21.7 °C (Table 7, Appendix A).

5.1.4 SANTA CLARA RIVER AT VCWPD PROJECT SIDE CHANNEL

The Santa Clara River at the Ventura County Watershed Protection District (VCWPD) Project side channel water temperature monitoring site was located in near the middle groin at the county bank stabilization project site off South Mountain Road in Santa Paula. The logger was wired to rocks on the bank and is resting on the bottom (Photo 11, Appendix B) in a pool about 3.5 ft deep. The objective at this location was to monitor water temperatures in the side channel of the Santa Clara River between Sespe Creek and the Freeman Diversion where resident trout captured in the downstream migrant trap are released during

the trapping season. In the future we will be adding more temperature monitoring sites throughout the Santa Clara River to closely monitor temperatures during the steelhead and Pacific lamprey migration season. The minimum water temperature collected at this site was 10.6 °C and the maximum temperature was 20.1 °C (Table 7, Appendix A).

5.1.5 SANTA PAULA CREEK UPSTREAM OF HARVEY DIVERSION

The Santa Paula Creek upstream of Harvey Diversion water temperature monitoring site was located in a low gradient riffle habitat type approximately 500 feet upstream from Harvey Diversion along Highway 150 in Santa Paula. The logger was attached to a fence post with zip ties under a boulder 0.4 inches from the bottom (Photo 12, Appendix B). The objective at this location was to monitor water temperatures in Santa Paula Creek. The minimum water temperature collected at this site was 6.8 °C and the maximum temperature was 27.1 °C (Table 7, Appendix A).

5.1.6 SANTA PAULA CREEK AT STECKEL PARK

The Santa Paula Creek at Steckel Park water temperature monitoring site was located in a pool habitat type ~50 feet upstream of the Steckel Park Bridge. The logger was attached 0.2 feet from the bottom with zip ties to a fence post. The fence post was located under a boulder and was secured to an alder tree via cable along the bank (Photo 13 and 14, Appendix B). This location was ideal for monitoring temperatures due to adequate flows, areas of scour and in-stream cover. However, the temperature logger was buried under sediment during a storm on March 18, 2011, but continued logging. This site is also currently isolated from the main flow. The minimum water temperature collected at this site was 10.0 °C and the maximum temperature was 20.9 °C (Table 7, Appendix A).

5.1.7 SANTA PAULA CREEK DIRECTLY DOWNSTREAM OF SISAR CREEK CONFLUENCE

This site was removed due to excessive tar accumulation on the logger and on the employees that accessed the site to upload data.

5.1.8 SESPE CREEK AT GRAND AVENUE

The Sespe Creek at Grand Avenue water temperature monitoring site was

located in a pool habitat type approximately 150 ft downstream of the USGS Gauge along Grand Avenue. The logger was attached to a fence post between two boulders, 0.5 feet from the bottom along the bank (Photo 15, Appendix B). The objective at this location was to monitor water temperatures at the resident and young-of-the-year coastal rainbow trout release location in Sespe Creek. This was an easily accessible pool with sufficient flow creating areas of scour and abundant in-stream cover, primarily riparian vegetation and boulder substrate. The minimum water temperature collected at this site was 8.3 °C and the maximum temperature was 30.7 °C (Table 7, Appendix A).

5.1.9 PIRU CREEK AT SANTA FELICIA DAM

The Piru Creek below Santa Felicia Dam water temperature monitoring site is located in a pool habitat approximately 500 ft downstream of the Santa Felicia Dam outlet works. Two data loggers were deployed at this site. The temperature logger (17C) was placed inside a standpipe above a pool water level logger (17B) which is at the deepest part of the pool (about 3 ft), 0.3 feet from the bottom along the bank (Photo 16, Appendix B). The objective at this location was to monitor water temperatures in the trout-bearing reach of Piru Creek below the dam. This was an easily accessible pool with sufficient flow creating areas of scour and abundant in-stream cover, primarily riparian and aquatic vegetation and boulders. The minimum water temperature collected at site 17B was 8.6 °C and the maximum temperature was 23.5 °C (Table 7, Appendix A). The minimum water temperature collected at site 17C was 8.7 °C and the maximum temperature was 23.4 °C (Table 7, Appendix A).

5.1.10 PIRU CREEK UPSTREAM OF SANTA CLARA RIVER CONFLUENCE

The Piru Creek upstream of the Santa Clara River confluence water temperature monitoring site is located in Piru Creek in a pool habitat approximately 1500 ft upstream of the Santa Clara River confluence. The data was taken from a water level logger inside a standpipe at the deepest part of the pool (about 2.0 ft), 0.2 feet from the bottom in the middle of the stream channel (Photo 17, Appendix B). The objective at this location was to monitor water temperatures near the terminus of Piru Creek where water temperatures can be very high in the summer months. The minimum water temperature collected at this site was 4.0 °C and the maximum temperature was 30.7 °C (Table 7, Appendix A).

5.4 DISCUSSION

The maximum water temperatures at most sites extends above what many believe to be the chronic or incipient upper lethal temperature limit of 25 °C for many anadromous salmonids, although higher temperatures reaching 29 °C can be tolerated for a short period of time (Myrick and Cech 2001) if water quality conditions are favorable and food is available. The maximum water temperature of 30.7 °C occurred in Sespe at Grand Avenue and Piru Creek at the Santa Clara River confluence during the summer. Both of these sites are near the terminus of each respective drainage in habitats lacking dense riparian vegetation. These extreme water temperatures are not uncommon in small streams of southern California. Southern California coastal rainbow trout have adapted to a wide variation in water temperatures by seeking out thermal refuges when available. Other behavioral responses to upper thermal extremes include increase feeding when food is available to offset the cost of an elevated metabolic rate although feeding does decline after about 19 °C but growth can still occur up to 25 °C (Myrick and Cech 2000). The duration of exposure is important because these fish will experience upper thermal limits over a period of a few hours during any given day but can maintain body weight at 25 °C for 30 days (Myrick and Cech 2000). In drainages where thermal refuge is limited or absent and low food production occurs, thermal stress alone can cause mortalities.

- Behler, J. L. and F. W. King. 1996. National Audubon Society Field Guide to North American Reptiles and Amphibians. Alfred A. Knopf, Inc. New York.
- California Department of Fish and Game. 2009. California Natural Diversity Database, Special Animals (883 taxa). The Natural Resources Agency, Biogeographic Data Branch.
- Chase, S. 2001. Contributions to the life history of adult Pacific lamprey (*Lampetra tridentata*) in the Santa Clara River of southern California. Bulletin of the Southern California Academy of Sciences 100(2):74–85.
- CMWD. 2008. 2008 Annual progress report for the Robles Diversion Fish Passage Facility. Casitas Municipal Water District, Oak View, CA.
- Haner, Philip V., J.C. Faler, R.M. Shrock, D.W. Rondorf, and A. G. Maule. 1995. Skin reflectance as a nonlethal measure of smoltification for juvenile salmonids. North American Journal of Fisheries Management 15: 814 - 822.
- Myrick, C. A., and J. J. Cech, Jr. 2000. Temperature influences on California rainbow trout physiological performance. Fish Physiology and Biochemistry 22(3):245–254.
- Myrick, C. A. and J. J. Cech, Jr. 2001. Bay-Delta Modeling Forum. Technical publication 01-1. Temperature effects on Chinook salmon and steelhead: a review focusing on California's Central Valley populations.
- National Marine Fisheries Service. 1997. Endangered and Threatened Species: Listing of Several Evolutionary Significant Units (ESUs) of West Coast Steelhead. Federal Register, 50 CFR Parts 222 and 227 [Docket No. 960730210–7193–02; I.D. 050294D] RIN 0648–XX65. Vol. 62, page 43937.
- National Marine Fisheries Service. 2003. Endangered and Threatened Species: Range Extension for Endangered Steelhead in Southern California. Federal Register, 50 CFR Part 224 [Docket No. 001025296-2079-02; I.D. 072600A] RIN 0648-AO05. Vol. 67 page 21586.
- National Marine Fisheries Service. 2006. Endangered and Threatened Species: Final Listing Determinations for 10 Distinct Population Segments of West Coast Steelhead. Federal

Register, 50 CFR Part 223 and 224 [Docket No. 051216341–5341–01; I.D. No. 052104F] RIN 0648–AR93. Vol. 71 page 834-862.

Rehmel, M. 2007. Application of Acoustic Doppler Velocimeters for Streamflow Measurements. *Journal of Hydraulic Engineering*. DOI: 10.1061/(ASCE)0733-9429(2007)133:12(1433).

Swift, C.C., T.R. Haglund, M. Ruiz, and R.N. Fisher. 1993. The status and distribution of the freshwater fishes of southern California. *Bull. S. Calif. Acad. Sci.* 92:101-167.

Swift, C. C., and S. R. Howard. 2009. Current Status and Distribution of the Pacific Lamprey South of Point Conception, Coastal Southern California, USA. *American Fisheries Society Symposium* 72:269-278.

Thomas R. Payne and Associates (TRPA). 2005. Assessing passage of steelhead over riffles in the lower Santa Clara River. Draft Final Report. 31 pp.

Wedemeyer, G.A. 1996. *Physiology of Fish in Intensive Culture Systems*. Chapman and Hall, New York, New York.

Appendix A

Tables and Figures

January																															
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
Ladder Operation																															
Smolt Trapping																															
Smolts																															
Adult Steelhead																															
Steelhead Kelt																															
Turbidity (ntu)	1.18	1.18	1.18	1.18	1.16	1.16	1.18	1.16	1.16	1.12	1.17	1.12	1.12	1.17	1.14	1.16	1.16	1.19	1.21	1.21	1.19	1.19	1.2	1.42	1.15	1.21	1.25	1.17	1.14	1.23	1.17
Estuary	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed
February																															
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28			
Ladder Operation																															
Smolt Trapping																															
Smolts																															
Adult Steelhead																															
Steelhead Kelt																															
Turbidity (ntu)	1.15	1.2	1.14	1.17	1.14	1.59	0.77	0.66	0.96	1.09	1.12	1.13	1.13	1.13	1.18	1.12	1.16	1.19	1.12	1.12	1.19	1.16	1.14	1.74	2.28	1.92	224	428			
Estuary	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Open	Open	Open	Open	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Open			
March																															
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
Ladder Operation																															
Smolt Trapping																															
Smolts																															
Adult Steelhead																															
Steelhead Kelt																															
Turbidity (ntu)	1729	1481	352	119	57	34.1	23.6	22.9	10.3	7.16	6.19	8.68	4.59	4.14	3.7	4.27	2.93	2.87	2.48	2.31	2.18	2.1	2.06	2.02	1.87	1.88	1.61	1.68	1.38	1.43	1.51
Estuary	Open	Open	Open	Open	Open	Open	Open	Open	Open	Open	Open	Open	Open	Open	Open	Open	Open	Open	Open	Open	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed
April																															
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	
Ladder Operation																															
Smolt Trapping																															
Smolts																															
Adult Steelhead																															
Steelhead Kelt																															
Turbidity (ntu)	2.07	3.64	1.85	1.63	1.65	1.45	1.33	1.27	1.57	1.96	2.2	2.16	2.17	2.35	2.36	2.57	1.97	2.47	2.02	1.97	1.84	1.75	1.65	1.77	1.74	1.77	1.43	1.4	1.6	1.61	
Estuary	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	
May																															
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
Ladder Operation																															
Smolt Trapping																															
Smolts																															
Adult Steelhead																															
Steelhead Kelt																															
Turbidity (ntu)	1.67	1.42	1.46	1.65	1.54	1.6	1.88	1.79	1.5	1.34	1.31	1.55	1.44	1.22	1.27	2.06	1.59	1.46	1.96	1.56	1.63	1.61	1.56	1.56	1.48	1.47	1.46	1.45	1.44	1.45	1.45
Estuary	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed	Closed
End of Monitoring Season																															

*Blank cells mean no activity or no data collected on that day

Table 2. Santa Clara River Fish Species

Common Name	Scientific Name	Status	Resident - Anadromous	Regulatory Status
Tidewater Goby	<i>Eucyclogobius newberryi</i>	Native	Resident	FE
Partially Armored Stickleback	<i>G. a. microcephalus</i>	Native	Resident	--
Unarmored Stickleback	<i>G.a. williamsoni</i>	Native	Resident	FE, SE, DFG: FP
Arroyo Chub	<i>Gila orcuttii</i>	Native	Resident	--
Santa Ana Sucker	<i>Catostomus santaanae</i>	Native*	Resident	*FT
Pacific Lamprey	<i>Entosphenus tridentatus</i>	Native	Anadromous	--
Rainbow Trout	<i>Oncorhynchus mykiss</i>	Native	Resident	--
Steelhead Trout	<i>Oncorhynchus mykiss</i>	Native	Anadromous	FE
Black Bullhead	<i>Ameiurus melas</i>	Introduced	Resident	--
Brown Bullhead	<i>Ameiurus nebulosus</i>	Introduced	Resident	--
Owens Sucker	<i>Catostomus fumeiventris</i>	Introduced	Resident	--
Hybrid Sucker	<i>C.santaanae x C. fumeiventris</i>	Introduced	Resident	--
Prickly Sculpin	<i>Cottus asper</i>	Introduced	Resident	--
Common Carp	<i>Cyprinus carpio</i>	Introduced	Resident	--
Threadfin Shad	<i>Dorosoma petenense</i>	Introduced	Resident	--
Mosquitofish	<i>Gambusia affinis</i>	Introduced	Resident	--
Channel Catfish	<i>Ictalurus punctatus</i>	Introduced	Resident	--
Green Sunfish	<i>Lepomis cyanellus</i>	Introduced	Resident	--
Bluegill	<i>Lepomis macrochirus</i>	Introduced	Resident	--
Mississippi Silverside	<i>Menidia audens</i>	Introduced	Resident	--
Largemouth Bass	<i>Micropterus salmoides</i>	Introduced	Resident	--
Fathead Minnow	<i>Pimephales promelas</i>	Introduced	Resident	--
Black Crappie	<i>Pomoxis nigromaculatus</i>	Introduced	Resident	--
White Crappie	<i>Pomoxis annularis</i>	Introduced	Resident	--
Brown Trout	<i>Salmo trutta trutta</i>	Introduced	Resident	--
Shimofuri goby	<i>Tridentiger bifasciatus</i>	Introduced	Resident	--

FE = Federally Endangered Species; FT = Federally Threatened Species; SE = State Endangered Species; DFG: FP = California Department of Fish & Game - Fully Protected Species.

*Santa Ana Sucker is not listed as FT in the Santa Clara River because it was considered introduced. Recent genetic analysis conducted by the USGS suggests that this species is native to the Santa Clara River.

Table 4 - Freeman Diversion Fish Monitoring Species Totals 2014

Fish Species: **STK** = Steelhead Adult Kelt, **ST** = Steelhead Adult, **RS** = Smolt, **RT** = Resident Rainbow, **YOY** = Young of the Year *O. mykiss*, **PL (Adult)** = Lamprey Adult, **PL (Am)** = Lamprey Ammocoete, **TS** = Stickleback, **AC** = Arroyo Chub, **SS** = Santa Ana Sucker, **OS** = Owens Sucker, **SSxOS** = Sucker Hybrid, **FM** = Fathead Minnow, **LB** = Largemouth Bass, **GS** = Green Sunfish, **BB** = Brown Bullhead, **BC** = Black Bullhead, **PS** = Prickly Sculpin, **MF** = Mosquitofish **CR** = Crappie spp., **SG** = Shimofuri Goby

Amphibian & Reptile Species: **WT*** = Western Toad, **AT*** = Arroyo Toad, **SFT*** = Spadefoot Toad, **BF** = Bullfrog, **CRLF** = Red-legged Frog, **PT*** = Pond Turtle, **AF** = African Clawed Frog, **TF*** = Tree Frog

Native species indicated with an asterisk*

Other: **RSC** = Red swamp crayfish

Total	Native Fishes									Non-native Fishes												
	STK	ST	RS	RT	YOY	PL (Adult)	PL (Am)	TS	AC	SS	OS	SS x OS	FM	LB	GS	BB	BC	CC	PS	MF	CR	SG
	0	0	11	4	4	0	0	608	20	6	17	4	57	0	6	1	13	10	52	4262	0	9

Total	Amphibians and Reptiles							Other	
	WT*	AT*	SFT*	CRLF*	BF	AF	TF*	PT*	RSC
	6	0	0	0	242	39	1	1	2813

Biologists: Mike Booth, Steve Howard **Fish Technicians:** Dave Wilson, Jessica Cornelius, Kelsey Erisman, Akiko Mandell

Table 5 - Santa Clara River Reptile and Amphibian Species

Common Name	Scientific Name	Status	Regulatory Status
Two-striped Garter Snake	<i>Thamnophis couchi hammondi</i>	Native	DFG: SSC
Southwestern Pond Turtle	<i>Clemmys marmorata pallida</i>	Native	DFG: SSC
Red-eared Slider	<i>Chrysemys scripta elegans</i>	Introduced	--
California Treefrog	<i>Hyla cadaverina</i>	Native	--
Pacific Treefrog	<i>Hyla regilla</i>	Native	--
Western Toad	<i>Bufo boreas</i>	Native	--
Bullfrog	<i>Rana catesbeiana</i>	Introduced	--
African Clawed Frog	<i>Xenopus laevis</i>	Introduced	--
DFG: SSC = California Department of Fish & Game - Species of Special Concern			

Table 6. Temperature Monitoring Sites

Location	Logger #	Date deployed	Max Depth	Logger depth (ft from bottom)	Habitat Type	Photo (Appendix B)
Freeman Fish Trap Bay	1269160	5/16/2008	3.0	1.0	pool	8
Freeman Fish Screen Bay	1269166	12/8/2008	2.5	0.5	pool	9
Santa Clara River at VCWPD Project	2270563	1/11/12	3.5	2.0	pool	10
Santa Clara River at VCWPD Project side channel	9947131	8/01/12	3.5	0.6	pool	11
Santa Paula Creek upstream of Harvey Diversion	2270574	1/13/12	0.9	0.3	riffle	12
Santa Paula Creek at Steckel Park	1269161	3/27/08	1.8	0.2	pool	13, 14
Santa Paula Creek directly downstream of Sisar Creek confluence	2250248	1/16/2009	Logger Pulled			
Sespe Creek at Grand Avenue	2270588	1/11/12	1.5	0.5	pool	15
Piru Creek at Santa Felicia Dam (level logger)	10185710	9/6/12	1.75	0.2	pool	16
Piru Creek at Santa Felicia Dam	2267160	11/6/12	1.75	0.2	pool	16
Piru Creek at Santa Clara River confluence	10185717	9/6/12	2.0	0.2	pool	17

Table 7. Water Temperature Statistics by Monitoring Site.

Location	Max Temp (°C)	Date Max Temp	Min Temp (°C)	Date Min Temp	Mean Temp (°C)
Freeman Fish Trap Bay	31.5*	9/15/14	1.72*	12/9/13	16.4
Freeman Fish Screen Bay	47.1*	9/16/14	-0.17*	12/8/13	3.59
Santa Clara River at VCWPD Project	21.7	5/15/14	10.4	3/2/14	16.8
Santa Clara River at VCWPD Project side channel	20.1	9/16/14	10.6	3/2/14	15.6
Santa Paula Creek upstream of Harvey Diversion	27.1	8/28/14	6.8	12/10/13	16.4
Santa Paula Creek at Steckel Park	20.9^	9/17/14	10.0^	12/13/13	15.9
Santa Paula Creek directly downstream of Sisar Creek confluence	Pulled Logger				
Sespe Creek at Grand Avenue	30.7	7/4/14	8.3	12/9/13	18.4
Piru Creek at Santa Felicia Dam (level logger)	23.4	9/16/14	8.7	1/1/14	15.8
Piru Creek at Santa Felicia Dam	23.5	9/16/14	8.6	1/18/14	15.7
Piru Creek at Santa Clara River confluence (level logger)	30.7	7/1/14	4.0	12/10/13	17.6

* Logger out of water.

^ Pool laterally isolated from creek, logger under rock.




Figure 1. Schematic of Freeman Diversion Facility

Figure 2 - Daily Rainfall Totals in Santa Paula, CA





Figure 3 - Steelhead Monitoring Results






Figure 4. Freeman Bypass Flow Monitoring Sites below the Freeman Diversion




Figure 5. Water Temperature Monitoring Sites within the Santa Clara River Watershed.

Figure 6. Freeman Fish Trap Bay Water Temperature




Figure 7. Freeman Fish Screen Bay Water Temperature




Figure 8. Santa Clara River at Ventura County Bank Stabilization Project




Figure 9. Santa Clara River Side Channel at Ventura County Bank Stabilization Project




Figure 10. Santa Paula Creek Upstream of Harvey Diversion




Figure 11. Santa Paula Creek at Steckel Park




Figure 12. Sespe Creek at Grand Avenue




Figure 13. Piru Creek at Santa Felicia Dam - Pool




Figure 14. Piru Creek at Santa Felicia Dam - Pool





Figure 15. Piru Creek Upstream of Santa Clara River Confluence - Pool



Appendix B

Photos



Photo 1. Santa Clara River estuary release site open



Photo 2 - Santa Clara River estuary release site closed



Photo 3. Santa Clara River steelhead smolt



Photo 4. Santa Clara River resident rainbow trout.



Photo 5. Upstream migration monitoring infrared scanners



Photo 6. Upstream migration monitoring surveillance system.



Photo 7. Fish screen bay stranding survey



Photo 8. Fish trap bay water temperature monitoring site



Photo 9. Fish screen bay water temperature monitoring site



Photo 10. VCWPD project water temperature monitoring site



Photo 11. VCWPD project side channel water temperature monitoring site



Photo 12. Santa Paula Creek upstream of Harvey Diversion water temperature monitoring site



Photo 13. Santa Paula Creek at Steckel Park water temperature monitoring site 2010 (this site has become overgrowth with riparian vegetation and has become isolated from the creek, See photo 14)



Photo 14. Santa Paula Creek at Steckel Park water temperature monitoring site 2014



Photo 15. Sespe Creek at Grand Avenue water temperature monitoring site



Photo 16. Piru Creek downstream of Santa Felicia Dam water temperature monitoring site



Photo 17. Piru Creek ~1500 feet upstream of Santa Clara River confluence water temperature monitoring site

Freeman Diversion Panoramic Photos



3/6/2014



3/13/2014



3/18/2014



3/24/2014



3/26/2014



3/30/2014



4/7/2014



4/13/2014



4/19/2014



4/25/2014



4/30/2014